VI. A Letter to the Learned Dr. Caleb Cotesworth, F. R. S. of the College of Physicians, London, and Physician to St. Thomas's Hospital; containing a Comparison between the Danger of the Natural Small Pox, and of that given by Inoculation. By James Jurin, M.D. R. S. Secret.

SIR,

The fincere and difinterested Regard for the Good of Mankind, which you have always manifested, as well in your extensive private Practice, as in that publick Post, which you have so long and so usefully sill'd, must affect you, I am sure, with a great and sensible Concern for the Destruction made among us by that terrible Calamity the Small Pox. We have seen, for some considerable Time past, above 100 Persons per Week in this City and Suburbs, taking one Week with another, to be carry'd off by this Disease; a Consideration certainly that ought to dispose us to enter into any Measures, by which we may reasonably hope to put some Stop to the Progress of so cruel a Dissemper.

To this Purpose, Sir, the Method of Inoculation, which has lately been introduced among us, is strongly recommended on the one hand, and has been opposed with a great deal of Warmth and Zeal on the other.

I have

I have no Inclination to enter into this Controverfy; it is in better and abler Hands: but, as the Point in Difpute is of the utmost Importance to Mankind, I heartily wish, that, without Passion, Prejudice, or private Views, it may be fairly and maturely examin'd. In order to which, if the following Extracts and Computations, concerning the comparative Danger of the Inoculated and Natural Small Pox, may be of any Use to your self, or to other impartial and disinterested Judges, I shall think my Labour well bestowed.

The Number of Persons, who have had the Small Pox by Inoculation here in *England*, is, by the best Information I have been able to collect, as follows.

Inoculated by	
Dr. Nettleton	-
Claudius Amyand, Esq; Serjeant Surgeon	61
Mr. Maitland, Surgeon	721.20
Dr. Dover	57
Mr. Weymish, Surgeon	- 4
The Reverend Mr. Johnson	3
Dr. Brady, at Port mouth	3
Mr. Smith, Surgeon, and Mr. Dymer, 2	4
Apothecary, at Chichester	13
Mr. Waller, Apothecary at Gosport	
A Woman at Leicester	3
Dr. Williams at Haverford Woa	8
Two other Persons near the same Place	6
Dr. French, at Bristol	- 2
The second of the second secon	I
In all	182

Out of this Number the Opposers of Inoculation affirm, that two Persons died of the Inoculated Small Pox; the Favourers of this Practice maintain, that their Death was occasioned by other Causes. If, to avoid dispute

dispure, these two be allow'd to have died of Inocularion, we must estimate the hazard of dying of the Inoculated Small Pox, as far as can be collected from our own Experience, to be that of 2 out of 182, or

one out of 91.

The Reverend Mr. Mather, in a Letter dated March 10. 1721. from Boston in New England, gives an Account, That of near 300 inoculated there, 5 or 6 died upon it or after it, but from other Diseases and Accidents, chiefly from having taken the Infection in the common way by Inspiration, before it could be given

them in this way of Transplantation.

If, as we have done before, to avoid all occasion of dispute, we allow 5 out of these 300 to have died of the Small Pox by Inoculation, notwithstanding what Mr. Mather has faid of their dying by other Accidents or Diseases; the hazard of Inoculation will thence be determin'd to be that of I in about 60. But here it must be observ'd, that by all the Accounts from New England, the Operators there appear not to have been so cautious in the choice of their Subjects. as here in England. For Mr. Mather tells us, that the Persons inoculated were young and old, from I Year to 70, weak and strong; and by other relations we are inform'd, that Women with Child, and others even in Childbed, underwent the Operation. Apparently the Greatness of the Danger they were in, from the Infection in the natural Way, which then raged among them with the utmost Fury, made them the more adventurous.

We come now, Sir, to the fecond Part of our Defign, which is to form an Estimate of the Hazard, which all Mankind, one with another, are under of dying of the natural Small Pox, that, by comparing this with the Hazard of Inoculation, the Publick may

M m 2

be enabled to form a Judgment, whether or no the Practice of Inoculation tends to the Preservation of Mankind, by lessening the Danger to which they are otherwise liable.

With this View I have confulted the Yearly Bills of Mortality, as far back as the Year 1667, being the Year after the Plague and the Fire of London, comprehending to the present Time the Space of 56 Years, from 42 of which I have given Extracts in the

two following Tables.

The first of these takes in the first 20 Years, distinguishing for every Year the total Number of Burials, and likewise the Number that died of the Small Pox, in two separate Columns. The third Column shews, how many died of the Small Pox out of every Thousand that were buried; and the fourth Column represents the Proportion between those that died of the Small Pox, and the whole Number of Burials, by the nearest Vulgar Fraction, having always 1 for the Numerator.

The second Table gives the last 22 Years, after the fame manner, and at the bottom of each Table is given the Total Number for each Series of Years, and likewise the Number that died each Year, taken at a Medium, one Year with another: by which it appears, that the Proportion between the Number of those that die of the Small Pox, and the whole Number of Burials, is very nearly the same, upon an Average for

each Series of Years.

The 14 intermediate Years between 1686 and 1701 are left out, because in the Bills for those Years, the Accounts of the Small Pox and Measles are not distinguished, as in the preceding and following Years, but are join'd together in one Article, fo that from them no certain Account can be drawn of the Number of Persons, that died of the Small Pox.

TABLE

## TABLE I.

es axof litus.	opinio had	La los	Total	
- normodera re	Total No	Died o	f the Sma	Il Pox.
Years	of Burials	In all.	In 1000	In Proportion
1667	15842	1196	75	33
1668	17278	1987	115	9
1669		951	49	20
1670	20198	1465	8.73	14
1671	15729	696	5 44	.23
1672	18230	1116	6 I	20.2
1673	17504	853	49	121
1674	21201	2507	58	2 2 27
1675	17244	997	SERVICE TO THE	I I V .
1676	18732	359 1678	88	132
1678	20678	1798	87	TT
1679	21730	1967	91	1 2
1680	21053	689	33	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1681	23971	2982	125	8
1682	20691	1408	68	75
1683	20587	2096	102	To
1684	23202	156	7	149
1685	23222	2496	107	30
1686	22609	1062	47	31
20 Years	398200	28459	71 1/2	¥ ¥4
			73.71	Course & C.S.
of Manager		HODE T	es into	he free l
Each Year at ? a Medium	19910	1423	71 1/2	74
a Meaium 5	tion is	o bil	unde lie	mr theur
Committee and	10 0872	8 650	90376	"As Vears
	TOTAL TI			
Call May Tella	WELL AND	Burke	are a	Endl Year in
CT I	50 - 72	21 16	igis .	42 at a 16-
		-		diam.

## TABLE II.

-	[7	Total Nº[	Died of the Small Pox.			
	Years				In Proporti	on.
-	1701	20471	1095	53	To	
	1702	19481	311	16	2.2 z	A
	1703	20720	898	43	24	
100	1704	22684	1501	66	H 10 H 10	
	1705	19847	721	50 36	20 1 28	
	1706	21600	1078	50	7 X	
	1707	21291	1687	79	73	
	1709	21800	1024	47	2 T	
	1710	24620	3138	127	3	
1111	1711	19833	915	46	22	. 1
	1712	21198	1943	92	TR	
	1713	21057	1614	77	Tã X	
	1714	26569	1057	48	2 2 2 2 2	No.
	1715	00	2427	99	E	
1	1717	23446	22 I I	94	TT	
1	1718	26523	1884	The second second	14	
	1719	28347	3229		9 2	
1	1720	25454	1440	57	18 11	
1	1721		2375	1	T Z	
1-	1722	-	36620		27537 6	77
1	22 Years	505598	30020	-	- 34	
1		ALL OF THE R	6.12	The state of		
I	Each Year at ?	22982	1665	72	74	
1	a Measum.	None and A	Sale	Taylor		
1	42 Years	903798	65079	72	<del>1</del>	7 9 1
1	42 Icars		-			
	Each Year in 7			7:	2 7 <sup>k</sup> / <sub>4</sub>	toling!
1	42 at a Me-	21519	15.5	1	14	
1		Pigli d	1		1 100	

By these Tables it appears, that upwards of seven per Cent. or somewhat more than a sourteenth part of Mankind, die of the Small Pon; and consequently the hazard of dying of that Distemper, to every individual born into the World, is at least that of one in sourteen. And that this Hazard encreases after the Birth, as the Child advances in Age, will appear from what follows.

From this Estimate it is demonstrable, that, in the Case of Persons actually having the Small Pox, the Hazard that they run, one with another, of dying of that Distemper, is greater than that of one in fourteen; or, which is the same thing, there must be fewer than thirteen, that recover, for one that dies of the Small Pox. For fince one fourteenth part of Mankind die of the Small Pox, and the other thirteen parts die of other Diseases; if these thirteen have all had the Small Pox, and recover'd from it, before they fell ill of those other Diseases of which they died, then just thirteen will have recover'd from the Small Pox, for one that dies of that Distemper: but, as it is notorious, that great Numbers, especially of young Children, die of other Diseases, without ever having the Small Pox, it is plain, that fewer than thirteen must recover from this Distemper, for one that dies of it.

To determine exactly how many of these thirteen Parts of Mankind, die without having the Small Pox, is a very difficult Task: but it is easy to see, that a considerable Deduction is to be made from them.

In the first place, the two Articles of Stilborn and Abortive Children, which are put into the yearly Bills, as part of the Number of Burials, are unquestionably to be deducted.

With these two, Sir, you will not I believe, think it unreasonable to join the following Heads, which, by the best Information I can procure, comprehend only very young Children, or at most not above one or two Years of Age. Overlaid, Chrysoms and Infants, Convulsions, Horseshoehead, Headmoldshot, Teeth, Water in the Head, Worms, Rickets, Livergrown, Chin-Cough, and Hooping-Cough, which Articles in the Yearly Bills for 22 Years last past, amount at a Medium to 386 in each 1000, of the whole Number of

It is true indeed, that, in all Probability, some fmall part of these must have gone thro' the Small Pox, and therefore ought not to be deducted out of the Account: but then, on the other hand, as it is certain,

that of the remaining  $\frac{614}{1000}$  of Mankind, that are above

one or two Years of Age, there are great Numbers, that never have the Small Pox, it will I presume be judged to be no unequal Supposition, if I suppose all that are contain'd under the Heads abovemention'd, to have miss'd that Distemper, when by way of Compensation, I allow all the remainder of Mankind to undergo it; which Concession is so large, that it will abundantly make up for what I assume too much in the former Supposition.

Allowing therefore, that out of every 1000 Children that are born, 386 die under one or two Years of Age, without having the Small Pox, and 72 do some time or other die of that Distemper; it follows, that the Hazard of dying of it, to the remainder of Mankind, above one or two Years of Age, who are all supposed to undergo that Disease sooner or later, is that of 72 out of 614, or nearly 2 out of 17: 10 that no more, than between 7 and 8, can recover from that Distemper, for one that dies of it. And if any consirable part of the aforesaid remainder of Mankind, more than is allow'd for above, do escape having the Small Pox, then the Proportion of those that recover from it, will be still smaller.

This Consideration shews the Fallacy of one plaufible Argument, that has been often made use of on occasion of the present Disputes about Inoculation: Which is, that whatever be the Danger of dying of the Small Pox, to those that actually have that Disease, yet, as great Numbers of Persons never have the Small Pox at all, this Danger is what any particular Person may never be in; and therefore it will be Madness to undergo the hazard of Inoculation, be it great or small, in order to prevent a Disease, which possibly may never

For if two parts in seventeen of all Mankind, that are above one or two Years of Age, must sooner or later die of the Small Pox, it is plain, that how many parts soever of these seventeen are supposed to escape that Distemper, the Mortality among the remainder, who undergo it, must in proportion be so much the greater. As for instance, if 7 parts escape having the Small Pox, and 10 undergo it, then 2 out of 10, or 1 out of 5, that have the Small Pox, must die of that Disease.

befal one.

And as it can never be known, whether or no any particular Person be one of those, that are to have the Small Pox; his Hazard of dying of that Distemper, being made up of the Hazard of having it, and the Hazard of dying of it, if he has it, will be exactly the same, namely, that of 2 in 17, or 1 in 8 or 9, whether the proportion of Mankind, that escape having the Small Pox, be great or small.

Nn But

But as what has been faid concerning the Hazard of the natural Small Pox, is taken from an Account of 42 Years; whereas the Hazard of Inoculation is estimated only from what has happen'd in the space of about 18 Months, since which time it had its first Rise among us; it will perhaps be asked by some Persons, why we do not likewise make the Estimate of the hazard of the natural Small Pox, from the last two Years alone, without running back into so great a number of Years, before Inoculation was begun?

To which we answer, that the Proportion of those that die of the Small Pox, varies so much in different Years, as appears from the Tables above, that it was impossible to come at any Certainty in this point, from the consideration of the two last Years alone: and if any one suspects us of Partiality in proceeding after the manner we have done, he need only cast his Eyes upon the second Table, where he will find, that the Mortality of the natural Small Pox, for the two last Years, has considerably exceeded the Medium we have determin'd, from taking in two and fourty Years.

There is another Method, which, if it were put in practice in several large Towns, or Parishes, and for a sufficient Number of Years, would enable us to come at a nearer and still more certain Computation of the Proportion between those that recover, and those that die of the Small Pox: which is, to send a careful Person once a Year, from house to house, to enquire what Persons have had the Small Pox, and how many have died of it, in the preceding Year. This has been done by Dr. Nettleton the last Year, at several Towns in Torkshire, &c. and the same was done at Chichester for the same Year, to the 15th of Ottober last, by a Person of Credit, whose Account was communicated to me by my learned and ingenious Friend, Dr. Whitaker.

Snch another Account has been transmitted to me from Haverford West, in South Wales, by the Learned Dr. Perrot Williams, Physician in that Place. The Sum of these Accounts is as follows.

Several Towns in Yorkshire	Sick of the Small Pox.	Died. 636
Chichester — — — Haverford West — —		168 52
he whole time of that Practi	Total 4626 —	856

From which it appears, that, upon a Medium between these Accounts, there died of the Small Pox almost 19 per Cent. or nearly one in five, of Persons of all Ages, that underwent that Distemper. Which is the more to be remarked, for that out of 82 Persons, that had the Small Pox by Inoculation, the same Year, and in the Neighbourhood of the same Places, not one miscarry'd.

Mr. Mather observes, in his Letter mention'd above, that out of more than 5000 Persons that had the Small Pox at Boston in New England, within little more than half a Year, near 900 died, which is more than one in six; and this Account added to those from Yorkshire, Chichester, and Wales, reduces the Proportion of those that die of the Small Pox to somewhat more than 18 per Cent. so that the Hazard of dying of that Distemper, to those who are taken ill of it, is that of one in between five and six, or something above two in eleven.

The Result therefore, Sir, of these Computations is, that, if the same Proportions should still continue, as have hitherto been determin'd by Observation, we must expect,

That

That of all the Children that are born, there will, fome time or other, die of the Small Pox, one in fourteen.

That of Persons of all Ages taken ill of the natural Small Pox, there will die of that Distemper, one in five

or six, or two in eleven.

That of Persons of all Ages inoculated, without regard to the Healthiness or Unhealthiness of the Subject, as was practifed in New England, there will die

one in sixty.

That of Persons inoculated with the same Caution in the choice of the Subjects, as has been used by the feveral Operators one with another, here in England, (if we allow in the two disputed Cases abovemention'd, that the Persons died of the inoculated Small Pox)

there will die one in ninety one.

But if those two Persons be allowed to have died of other Accidents or Diseases, then we shall have reason to think, as far as any Judgement can be made from our own Experience here in England, that none at all will die of Inoculation, provided that proper Caution be used; as we are inform'd, on all hands, is the Case in Turkey: where out of many thousands, that, in the space of about fourty Years past, have been inoculated in and about Constantinople, by one Greek Woman, who still continues that Practice, notwithstanding her extreme old Age, not so much as one Person has miscarried, as I am affured by the ingenious Dr. le Duc, a Native of Constantinople, who was himself inoculated there under the Care of his Father, an eminent Physician in that City. that, of the lame Proportions thould this

N. n. 2

I am, &c. do vd b'niscreto been determin'd by Ob . 28.